

New Combinations in *Urceola* (Apocynaceae)

David J. Middleton

Department of Botany, Trinity College, University of Dublin, Dublin, Ireland

ABSTRACT. Seven new combinations in *Urceola*, *U. xylinabariopsoides*, *U. huaitingii*, *U. micrantha*, *U. quintaretii*, *U. tournieri*, *U. rosea*, and *U. linearicarpa*, are proposed.

In a forthcoming paper in *Kew Bulletin*, the relationships between *Urceola* Roxburgh, *Ecdysanthera* Hooker & Arnott, *Parabarium* Pierre ex Spire, *Xylinabaria* Pierre, and *Xylinabariopsis* Pitard will be discussed, and the conclusion reached that these genera are not distinguishable (Middleton, in press). A revision of *Urceola* in Thailand, with the generic name *Urceola* taking priority, will be presented.

The problem now arises that there are no combinations in *Urceola* for those species of *Ecdysanthera* that are to appear in the account of the Apocynaceae for the *Flora of China*. This paper seeks to rectify this situation until such a time as the entire genus can be satisfactorily revised.

***Urceola xylinabariopsoides* (Tsiang) D. J. Middleton, comb. nov.** Basionym: *Chunechites xylinabariopsoides* Tsiang, Sunyatsenia 3: 306. 1937. TYPE: China. Hainan: Lok-tung, 9 June 1936, S. K. Lau 27035 (holotype, SYS).

***Urceola huaitingii* (Chun & Tsiang) D. J. Middleton, comb. nov.** Basionym: *Parabarium huaitingii* Chun & Tsiang, J. Arnold Arbor. 28: 245. 1947. TYPE: China. Guangxi (as Kwangsi): Shih-Wan-Ta Shan, Hsiang-Tze, 30 Apr. 1944, S. H. Chun 5027 (holotype, SYS).

***Urceola micrantha* (Wallich ex G. Don) D. J. Middleton, comb. nov.** Basionym: *Echites mi-*

crantha Wallich ex G. Don, Gen. Hist. 4: 75. 1837. TYPE: India. Pundua, *Wallich* 1667 (lectotype, selected by Lý (1978), CGE).

***Urceola quintaretii* (Pierre) D. J. Middleton, comb. nov.** Basionym: *Ecdysanthera quintaretii* Pierre, Rev. Cult. Colon. 11: 228. 1902. TYPE: Laos. Camon, *Tournier?* 6618 (lectotype, selected by Lý (1978), P).

***Urceola tournieri* (Pierre) D. J. Middleton, comb. nov.** Basionym: *Ecdysanthera tournieri* Pierre, Rev. Cult. Colon. 11: 228. 1902. TYPE: N Laos. Tran-ninh, *Tournier* 6613 (lectotype, selected here, P).

***Urceola rosea* (Hooker & Arnott) D. J. Middleton, comb. nov.** Basionym: *Ecdysanthera rosea* Hooker & Arnott, Bot. Beechy Voy. 198. 1836. TYPE: China. *Vachel* 144 (lectotype, selected here, K).

***Urceola linearicarpa* (Pierre) D. J. Middleton, comb. nov.** Basionym: *Ecdysanthera linearicarpa* Pierre, Rev. Cult. Colon. 11: 228. 1902. TYPE: N Laos. Tran-ninh, *Tournier* 6615 (lectotype, selected by Lý (1978), P).

Acknowledgment. I am grateful to Ihsan Al-Shehbaz for his help in preparing this paper.

Literature Cited

- Lý, T. D. 1978. Beiträge zur Kenntnis der Sippenstruktur des Genus *Parabarium* Pierre ex Spire (Apocynaceae). Feddes Repert. 89: 205–292.
Middleton, D. A revision of *Urceola* (Apocynaceae) in Thailand. Kew Bull. 49(4) (in press).